

Floramite™ SC

Miticide

Selective miticide for the control of two-spotted spider mites and Lewis mites on indoor ornamentals grown in greenhouses, shadehouses and interiorscapes. For the control of two-spotted spider mites on greenhouse vegetables (tomatoes, cucumbers, peppers); and for the control of spruce spider mite in Outdoor Grown Conifers including Christmas trees.

COMMERCIAL

GUARANTEE

bifenazate: 22.6 %

Contains 1,2-benzisothiazolin-3-one at 0.01% as a preservative

REGISTRATION NO: 27924
PEST CONTROL PRODUCTS ACT

READ THE LABEL BEFORE USING

KEEP OUT OF REACH OF CHILDREN

CAUTION – EYE IRRITANT



NET CONTENTS: 946 mL

**Chemtura Canada
Co./Cie**

25 Erb Street
Elmira, ON N3B 3A3

**☎ 24 Hour Emergency Phone:
1-866-744-3060**

**Grower Information:
1-800-350-1745**

™Floramite is a Trademark of Chemtura Canada Co./Cie

GENERAL INFORMATION:

Floramite SC Miticide is a selective miticide for the control of two-spotted spider mites and Lewis mites on indoor ornamentals grown in greenhouses, shadehouses and interiorscapes. Floramite SC Miticide also controls two-spotted spider mites on greenhouse vegetables (tomato, cucumber, pepper); and spruce spider mites on outdoor grown conifers including Christmas trees (i.e., spruce, pine, fir and cypress).

When used as directed and applied to the foliage, Floramite SC Miticide provides quick knockdown through contact activity.

DIRECTIONS FOR USE:**Mixing Instructions:**

Fill the spray tank with 1/2 the desired amount of water. Then add the required amount of Floramite SC with agitation running to fully disperse the product. Then fill the tank with the remaining amount of required water.

Floramite SC has been shown to degrade rapidly when mixed and stored with alkaline water of high temperature (50° C). To prevent degradation under alkaline conditions, solutions of Floramite SC must be used promptly. A commercially available buffering adjuvant should be added to the solution to reduce the pH to a neutral/acidic range.

USE RATES AND RECOMMENDATIONS:

Ornamentals and greenhouse vegetables: Mix 133 mls Floramite SC Miticide (30 g ai) in 400 L water and apply as a full coverage spray to the foliage to obtain uniform coverage. Actual spray volume will vary depending on the size of the plants being treated. Application should be made as soon as mites appear and will provide residual control for up to 28 days.

Crop	Mites Controlled	Amount of Floramite SC Miticide per 400L	Total Number of Sprays per Crop Cycle	Restricted Entry Interval (REI)
Indoor ornamentals grown in greenhouses, shadehouse and interiorscapes	Two-Spotted Spider Mite Lewis Mite	133 ml	1	12 hours

Crop	Mites Controlled	Amount of Floramite SC Miticide per 400L	Total Number of Sprays per Crop Cycle	Restricted Entry Interval (REI)	Pre Harvest Interval (PHI)
Greenhouse Tomatoes(*)	Two-Spotted Spider Mite	133 mL	2	12 hours	0 days
Greenhouse Cucumbers(*) Peppers(*)					1 day

(*) Make only one application of Floramite SC Miticide before rotating to products of an alternative chemical class

Note: It is not effective against rust mites, broad mites and flat mites. Floramite SC Miticide is primarily active on the motile stages of mites.

Outdoor Grown Conifers:

Mix 625 mL Floramite SC Miticide (150 g a.i.) in 1000L water and apply as a full coverage spray to the foliage to obtain uniform coverage. Actual spray volume will vary depending on the size of the plants being treated. Application should be made as soon as mites appear and will provide residual control for up to 28 days.

Crop	Mites Controlled	Amount of Floramite SC Miticide per 1000L	Number of Sprays per Year	Restricted Entry Interval (REI)
Outdoor Grown Conifers including Christmas trees (i.e. spruce, pine, fir and cypress)	Spruce Spider Mite	625 mL Note: It is important to keep the concentration of bifenthrin (i.e., 625 mL Floramite SC Miticide / 1000 L of water) constant if using a higher water volume (i.e., 625 mL Floramite SC Miticide in 1000 L/ha is equivalent to 1250 mL Floramite SC Miticide in 2000 L/ha).	Up to 2 applications/year. Use in sufficient water to obtain uniform coverage.	12 hours

Restrictions: Do not use Floramite SC Miticide more than twice per year. Do not use Floramite SC Miticide in successive applications. Apply only one application of Floramite SC before rotating to products of an alternative chemical class.

STRATEGIES FOR RESISTANCE MANAGEMENT

For resistance management, please note that Floramite SC Miticide contains a Group 25 acaricide. Any mite population may contain individuals naturally resistant to Floramite SC Miticide and other Group 25 acaricides. The resistant individuals may dominate the mite population if this group of acaricides is used repeatedly in the same greenhouses, shadehouses and interiorscapes. Other resistance mechanisms that are not linked to site of action but are specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay acaricide resistance:

- Ensure thorough spray coverage to all foliage.
- Where possible, rotate the use of Floramite SC Miticide or other Group 25 acaricides with different groups that control the same pests in a greenhouse, shadehouse or interiorscape.
- Acaricide use should be based on an IPM program that includes scouting, record keeping, and considers cultural, biological and other chemical control practices.
- Monitor treated pest populations for resistance development.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pest problems in your area.

Plant Tolerance

Neither the manufacturer, or the seller has determined whether or not Floramite SC Miticide can be used safely on all ornamental and greenhouse vegetable plants. Floramite SC Miticide has been tested on a variety of ornamental plants with no phytotoxicity observed at the label rate. However, all plant species and their varieties and cultivars have not been tested with sequential pesticide treatments. Therefore, prior to any large-scale application to ornamental plants or greenhouse vegetable plants, the user should determine the safety of Floramite SC Miticide by testing a small number of plant cultivars to be treated at the recommended rates to ensure that phytotoxic response will not occur. The end-user assumes all risks arising out of application in a manner inconsistent with this label.

PRECAUTIONS:

1. KEEP OUT OF REACH OF CHILDREN.
2. May cause moderate eye irritation. Avoid contact with eyes or clothing.
3. During mixing, loading, clean-up and repair workers MUST WEAR gloves, long-sleeved shirt, long pants and shoes plus socks. During application, workers MUST WEAR long-sleeved shirt, long pants and shoes plus socks.
4. If pesticide is spilled on work clothing, remove immediately. Wash thoroughly and put on clean clothing.
5. Work clothing should be washed with detergent and hot water. Wash work clothing separately from other laundry.
6. Do NOT contaminate food or feed products.
7. If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., consult CropLife Canada's website at www.croplife.ca.

ENVIRONMENTAL HAZARDS:

This product is toxic to aquatic organisms. Do not discharge effluent, waste and drainage water containing this product into water bodies, such as lakes, streams, ponds, rivers, and estuaries.

FIRST AID:

IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give **any** liquid to the person. Do not give anything by mouth to an unconscious person.

IF IN EYES: Hold eyes open and rinse slowly and gently with water for 15 – 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 – 20 minutes. Call a poison control centre or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION: Treat symptomatically.

STORAGE AND DISPOSAL:

STORAGE: Store product in original container only, away from other pesticides, fertilizer, food or feed.

DISPOSAL:

Do not reuse this container for any other purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple or pressure rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.
3. If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For information on the disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER:

This control product is to be used only in accordance with the directions on this label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of the product.

ID# 27924 – 080806

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.