



GROUP	9	FUNGICIDE
-------	---	-----------

**NET CONTENTS:  
2, 10 LITRES to BULK**

SCALA® SC FUNGICIDE

For control of listed leaf and fruit diseases in apples, pears, grapes, strawberries, potatoes and greenhouse tomatoes

COMMERCIAL

SUSPENSION



CAUTION

POISON



CAUTION – SKIN IRRITANT

REGISTRATION NUMBER: 28011 PEST CONTROL PRODUCTS ACT

GUARANTEE:

Pyrimethanil 400 g/L

Contains 1, 2-benzisothiazolin-3-one at 0.017% as a preservative.  
Warning, contains the allergen sulfites.

READ THE LABEL AND BOOKLET BEFORE USING  
KEEP OUT OF REACH OF CHILDREN

Notification Change

**For product information, call: 1-888-283-6847**

Bayer CropScience Inc.  
~~#100-3131, 114<sup>th</sup> Avenue SE~~  
~~Calgary, AB T2Z 3X2~~

Suite 200, 160 Quarry Park Blvd. S.E.  
Calgary, Alberta T2C 3G3

In case of spills, poisoning or fire, telephone emergency response number 1-800-334-7577 (24 hours a day).

**PRECAUTIONS:**

- KEEP OUT OF REACH OF CHILDREN
- Harmful if inhaled. Avoid inhaling/breathing sprays.
- May irritate the skin. Avoid contact with skin, eyes and clothing.
- When applied as a tank-mix combination, read and observe all label directions, including rates, restrictions, and grazing limitations for each product used in the tank-mix. Follow

the more stringent label precautionary measures for mixing, loading and applying stated on both labels.

- Apply only when the potential for drift to areas of human habitation or areas of human activity such as houses, cottages, schools, and recreation areas is minimal. Take into consideration wind speed, wind direction, temperature, application equipment and sprayer settings.
- Wash thoroughly with soap and water after handling and before eating, drinking, or smoking.
- If product comes in contact with clothing, remove all contaminated clothing, wash skin with soap and water and dress in clean clothing. Launder applicator clothing separate from other laundry.
- If this pest control product is to be used on a commodity that may be exported to the United States and you require information on acceptable residue levels in the United States, visit CropLife Canada's web site at [www.croplife.ca](http://www.croplife.ca).

#### **ENVIRONMENTAL HAZARDS:**

- Toxic to aquatic organisms. Observe buffer zones specified under DIRECTIONS FOR USE.
- To reduce runoff from treated area into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil, or clay.
- Avoid application of this product when heavy rain is forecast.
- Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.
- DO NOT apply this product through any type of irrigation system.
- DO NOT allow effluent or runoff from greenhouses containing this product to enter lakes, streams, ponds or other waters.

#### **PROTECTIVE CLOTHING AND EQUIPMENT:**

- During mixing, loading, clean-up and repair activities, workers must wear long pants, long-sleeved shirts, boots, chemical resistant gloves and safety goggles or a face shield.
- Applicators must wear long pants, long-sleeved shirts and boots.

#### **FIRST AID:**

**Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.**

**IF SWALLOWED:** Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

**IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

**IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

**TOXICOLOGICAL INFORMATION:**

- Treat symptomatically. Medical Personnel should contact Bayer's Medical Information Services, Toll-Free: 1-800-334-7577.

**STORAGE:**

- Do not contaminate water, food, or feed by storage or disposal.
- CANNOT BE STORED BELOW FREEZING.
- If stored for 1 year or longer, shake well before using.
- Store the tightly closed container away from feeds, seeds, fertilizer, plants and foodstuffs.
- Do not use or store in or around the home.
- Keep the product in the original container during storage.
- In case of fire, leaking or damaged containers, call toll free 1-800-334-7577.

**DISPOSAL:****Recyclable Container Disposal:**

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

**Disposal of Unused, Unwanted Product:**

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for cleanup of spills.

**NOTICE TO USER:**

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

SCALA is a registered trademark of Bayer.

091005



GROUP	9	FUNGICIDE
-------	---	-----------

**NET CONTENTS:  
2, 10 LITRES to BULK**

**SCALA® SC FUNGICIDE**

For control of listed leaf and fruit diseases in apples, pears, grapes, strawberries, potatoes and greenhouse tomatoes

COMMERCIAL

SUSPENSION



CAUTION

POISON



CAUTION – SKIN IRRITANT

REGISTRATION NUMBER: 28011 PEST CONTROL PRODUCTS ACT

GUARANTEE:

Pyrimethanil 400 g/L

Contains 1, 2-benzisothiazolin-3-one at 0.017% as a preservative.  
Warning, contains the allergen sulfites.

READ THE LABEL AND BOOKLET BEFORE USING  
KEEP OUT OF REACH OF CHILDREN

**For product information, call: 1-888-283-6847**

Notification Change

Bayer CropScience Inc.  
#100-3131, 114<sup>th</sup> Avenue SE  
Calgary, AB T2Z 3X2

Suite 200, 160 Quarry Park Blvd. S.E.  
Calgary, Alberta T2C 3G3

In case of spills, poisoning or fire, telephone emergency response number 1-800-334-7577 (24 hours a day)

# SCALA® SC FUNGICIDE

## Table of Contents:

## Section Number:

GENERAL INFORMATION	The Product	1
SAFETY AND HANDLING	Precautions, Protective Clothing & Equipment, and Re-entry Restriction	2
	First Aid and Toxicological Information	3
	Environmental Hazards	4
	Storage	5
	Disposal	6
	DIRECTIONS FOR USE	Application Information and Precautions
Crop, Disease, Rate, Timing, Minor Uses		8
Resistance Management		9
Mixing Instructions		10
Preharvest Intervals		11
Notices		12

FOR MORE INFORMATION CONTACT:

Notification Change

**Bayer CropScience Inc.**  
~~#100-3131-114<sup>th</sup> Avenue SE~~  
~~Calgary, AB T2Z 3X2~~

Suite 200, 160 Quarry Park Blvd. S.E.  
Calgary, Alberta T2C 3G3

## GENERAL INFORMATION

### Section 1: The Product

#### THE PRODUCT:

- For control of Botrytis bunch rot in grapes, Botrytis grey mould in strawberries, scab in apples and pears, early blight in potatoes and *Botrytis* storage diseases on apples, and suppression of *Penicillium* storage diseases on apples.
- SCALA® SC FUNGICIDE is a systemic fungicide that is effective on the above noted diseases. SCALA SC FUNGICIDE is best suited for use in a preventative treatment program.

## **SAFETY AND HANDLING**

### **Section 2: Precautions, Protective Clothing and Equipment, and Re-entry Restriction.**

#### **PRECAUTIONS:**

- KEEP OUT OF REACH OF CHILDREN
- Harmful if inhaled. Avoid inhaling/breathing sprays.
- May irritate the skin. Avoid contact with skin, eyes and clothing.
- When applied as a tank-mix combination, read and observe all label directions, including rates, restrictions, and grazing limitations for each product used in the tank-mix. Follow the more stringent label precautionary measures for mixing, loading and applying stated on both labels.
- Apply only when the potential for drift to areas of human habitation or areas of human activity such as houses, cottages, schools, and recreation areas is minimal. Take into consideration wind speed, wind direction, temperature, application equipment and sprayer settings.
- Wash thoroughly with soap and water after handling and before eating, drinking, or smoking.
- If product comes in contact with clothing, remove all contaminated clothing, wash skin with soap and water and dress in clean clothing. Launder applicator clothing separate from other laundry.
- If this pest control product is to be used on a commodity that may be exported to the United States and you require information on acceptable residue levels in the United States, visit CropLife Canada's web site at [www.croplife.ca](http://www.croplife.ca) .

#### **PROTECTIVE CLOTHING AND EQUIPMENT:**

- During mixing, loading, clean-up and repair activities, workers must wear long pants, long-sleeved shirts, boots, chemical resistant gloves and safety goggles or a face shield.
- Applicators must wear long pants, long-sleeved shirts and boots.

#### **RE-ENTRY RESTRICTION:**

- For apples and pears, DO NOT enter, or allow worker reentry, within 24 hours after application to carry out hand thinning activities.
- For grapes, DO NOT enter, or allow worker entry, within 24 hours after application to carry out cane turning, girdling, training, tying, pruning, thinning and leaf pulling activities.

### **Section 3: First Aid and Toxicological Information**

#### **FIRST AID:**

**Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.**

**IF SWALLOWED:** Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

**IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

**IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

#### **TOXICOLOGICAL INFORMATION:**

- Treat symptomatically. Medical Personnel should contact Bayer's Medical Information Services, Toll-Free: 1-800-334-7577.

#### **Section 4: Environmental Hazards**

##### **ENVIRONMENTAL HAZARDS:**

- TOXIC to aquatic organisms. Observe buffer zones specified under DIRECTIONS FOR USE.
- To reduce runoff from treated area into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil, or clay.
- Avoid application of this product when heavy rain is forecast.
- Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.
- DO NOT apply this product through any type of irrigation system.
- DO NOT allow effluent or runoff from greenhouses containing this product to enter lakes, streams, ponds or other waters.

#### **Section 5: Storage**

##### **STORAGE:**

- Do not contaminate water, food, or feed by storage or disposal.
- CANNOT BE STORED BELOW FREEZING.
- If stored for 1 year or longer, shake well before using.
- Store the tightly closed container away from feeds, seeds, fertilizer, plants and foodstuffs.
- Do not use or store in or around the home.
- Keep the product in the original container during storage.
- In case of fire, leaking or damaged containers, call toll free 1-800-334-7577.

#### **Section 6: Disposal**

##### **Recyclable Container Disposal:**

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed

of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

#### **Disposal of Unused, Unwanted Product:**

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for cleanup of spills.

#### **Disposal of Unused Spray Solution:**

If any spray solution remains in the tank after spraying is finished, it should be sprayed on the perimeter of the area just sprayed, away from water supplies, ditches, and irrigation canals.

## **DIRECTIONS FOR USE**

### **Section 7: Application Information and Precautions**

#### **APPLICATION INFORMATION:**

Thorough coverage of all plant parts to be protected is essential for good disease control. Use sufficient water volume and spray pressure to provide thorough and uniform coverage for optimum disease control.

Fruit Crops – Apply SCALA SC FUNGICIDE in a minimum spray volume of 1000 L/ha.

Vegetable Crops – Apply SCALA SC FUNGICIDE in a minimum spray volume of 300 L/ha for ground application and a minimum of 35 L/ha for aerial application.

Greenhouse tomatoes - Apply SCALA SC FUNGICIDE in a minimum spray volume of 250-600 L/ha.

#### **APPLICATION PRECAUTIONS:**

DO NOT apply more than 6.0 L/ha of SCALA SC FUNGICIDE in a single growing season. As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.

Field sprayer application: **DO NOT** apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE) medium classification. Boom height must be 60 cm or less above the crop or ground.

Airblast application: **DO NOT** apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** direct spray above plants to be treated. Turn off outward pointing nozzles at row ends and outer rows. **DO NOT** apply when

wind speed is greater than 16 km/h at the application site as measured outside of the treatment area on the upwind side.

Aerial Application: **DO NOT** apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply when wind speed is greater than 16 km/h at flying height at the site of application. **DO NOT** apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE) medium classification. To reduce drift caused by turbulent wingtip vortices, the nozzle distribution along the spray boom length **MUST NOT** exceed 65% of the wing- or rotorspan.

Apply SCALA SC FUNGICIDE only by rotary or fixed-wing aircraft which has been functionally and operationally calibrated for the atmospheric conditions of the area and the application rates and conditions of this label.

Label rates, conditions and precautions are product specific. Read and understand the entire label before opening this product. Apply only at the rate recommended for aerial application on this label. Where no rate for aerial application appears for the specific use, this product cannot be applied by any type of aerial equipment.

Ensure uniform application. To avoid streaked, uneven or overlapped application, use appropriate marking devices.

Use Precautions for Aerial Applications:

Apply only when meteorological conditions at the treatment site allow for complete and even crop coverage. Apply only under conditions of good practice specific to aerial application as outlined in the National Aerial Pesticide Application Manual, developed by the Federal/Provincial/Territorial Committee on Pest Management and Pesticides.

Apply only when the potential for drift to areas of human habitation or areas of human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature, application equipment and sprayer settings.

Operator Precautions for Aerial Applications:

Do not allow the pilot to mix chemicals to be loaded onto the aircraft. Loading of premixed chemicals with a closed system is permitted.

It is desirable that the pilot has communication capabilities at each treatment site at the time of application.

The field crew and the mixer/loaders must wear chemical resistant gloves, coveralls and goggles or face shield during mixing/loading, cleanup and repair. Follow the more stringent label precautions in cases where the operator precautions exceed the generic label recommendations on the existing ground boom label. All personnel on the job site must wash hands and face thoroughly before eating and drinking. Protective clothing, aircraft cockpit and vehicle cabs must be decontaminated regularly.

Product Specific Precautions for Aerial Applications:

Read and understand the entire label before opening this product. If you have questions, call the manufacturer at 1-888-238-6847 or obtain technical advice from the distributor or your provincial agricultural representative.

**Buffer zones:**

Use of the following spray methods or equipment **DO NOT** require a buffer zone: hand-held or backpack sprayer and spot treatment.

The buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands).

Method of application	Crop		Buffer Zones (metres) Required for the Protection of Freshwater Habitat of Depths:	
			Less than 1 m	Greater than 1 m
Field sprayer	Potato, strawberry		1	1
Airblast	Apple, pear	Early growth stage	15	1
		Late growth stage	5	1
	Grape	Early growth stage	20	1
		Late growth stage	10	1
Aerial	Potato	Fixed and rotary wing	10	1

For tank mixes, consult the labels of the tank-mix partners and observe the largest (most restrictive) buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank mix partners.

**ROTATIONAL CROP RESTRICTIONS:**

Wheat may be planted 30 days following the application of SCALA SC FUNGICIDE. For all other crops not listed on the SCALA SC FUNGICIDE label, a plant-back interval of 130 days must be observed.

**Section 8: Crop, Disease, Rate, Timing**

CROP	DISEASE	RATE	TIMING
Apple Pear	Scab	<u>SCALA SC FUNGICIDE</u> applied alone. Apply SCALA SC FUNGICIDE at 0.75 to 1.0 L/ha. Use the higher rate under high disease pressures. Ensure this rate is maintained in concentrate or dilute spray applications. Apply a maximum of 4 applications for scab alone.	Apply early in the season at 7-12 day intervals, from green tip to petal drop. For scab control, do not apply post-bloom. PHI of 72 days for apple and pear.  Apply by ground only.

Apple	Storage diseases:  Control of Botrytis  Suppression of Penicillium	<u>SCALA SC FUNGICIDE applied alone.</u> <b>FOR USE IN ONTARIO AND QUEBEC ONLY.</b> Apply a single application of SCALA SC FUNGICIDE at 1.5 to 2.0 L/ha. Use the higher rate if there is a history of high disease pressures in the field or storage facility.	Apply 2 weeks before harvest. PHI of 14 days.  Apply by ground only.
Grape	Botrytis Bunch Rot	<u>SCALA SC FUNGICIDE applied alone.</u> Apply SCALA SC FUNGICIDE at 2.0 L/ha. Apply maximum of 3 applications. Ensure resistance management recommendations for <i>Botrytis</i> are followed in Section 9.	Begin applications at early bloom if needed then follow with applications from berry touch to bunch closure. An application can also be made from fruit ripening through to harvest during frequent wet conditions (conducive to infection) or when symptoms of infection are evident. Thorough coverage of bunches is essential. Applications must be made a minimum of 7 days apart. PHI of 7 days.  Apply by ground only.
Strawberry	Botrytis Grey Mould	<u>SCALA SC FUNGICIDE applied alone.</u> Apply SCALA SC FUNGICIDE at 2.0 L/ha. Apply maximum of 3 applications. Ensure resistance management recommendations for <i>Botrytis</i> are followed in Section 9.	Make first application at the white bud stage (prebloom) and repeat applications as required at 7-10 day intervals. PHI of 1 day.  Apply by ground only.
Potato	Early Blight	<u>SCALA SC FUNGICIDE applied in tank mixture with BRAVO 500.</u> Apply SCALA SC FUNGICIDE at 750 mL/ha plus BRAVO 500 according to BRAVO 500 label directions. A maximum of six applications per year may be made.	Apply when plants are 15-20 cm high or when environmental conditions are conducive to disease development. Repeat applications at 7-14 day intervals or as necessary to maintain disease control. If severe disease conditions exist, use the 7 day interval. Minimum spray interval is 7 days. Ensure that the area to be treated is covered uniformly. PHI of 7 days.  Apply by ground or air.

## MINOR USES

NOTE TO USER: READ THE FOLLOWING BEFORE USING THIS PRODUCT FOR THE INDICATED SPECIAL USE APPLICATIONS (BELOW): The DIRECTIONS FOR USE for this product for the uses described below were developed by persons other than Bayer CropScience and accepted for registration by Health Canada under the User Requested Minor Use Label Expansion program. Bayer CropScience itself makes no representation or warranty with respect to performance (efficacy) or crop tolerance (phytotoxicity) claims for this product when used on the crops listed below. Accordingly, the Buyer and User assume all risks related to performance and crop tolerance arising, and agree to hold Bayer CropScience harmless from any claims based on efficacy or phytotoxicity in connection with the uses described below.

CROP	DISEASE	RATE	TIMING
Greenhouse tomatoes	7-day control of  Grey mould ( <i>Botrytis cinerea</i> )	2 L/ha	Apply Scala once only at first sign of Grey mould or as a rotational fungicide with other registered Grey mould control products in a spray program. Scala will provide control of Grey mould for 7 days after application depending on disease conditions. Ensure that the tomato crop is covered uniformly by the spray treatment.

RESTRICTIONS: Apply SCALA SC only in well-ventilated plastic tunnel houses or glass houses. Ventilate for at least 2 hours after application. Do not apply more than 1 application of SCALA SC per growing season to greenhouse tomatoes. Do not apply within 1 day of harvest. To limit the potential for resistance to develop, DO NOT apply more than one application of SCALA SC or other Group 9 fungicide before alternating with a fungicide from a different resistance management group.

Notice: In ventilated glass or plastic houses, the vapour activity of SCALA SC can produce brown or necrotic spots on the crop. Intensity and frequency of these effects depend on the relative humidity (e.g., above 80%), duration of vapour exposure and concentration of the spray solution. Thus, proper ventilation after spraying is essential in greenhouse applications.

## Section 9: Resistance Management

### RESISTANCE MANAGEMENT RECOMMENDATIONS:

For resistance management, please note that SCALA SC FUNGICIDE contains a Group 9 fungicide. Any fungal population may contain individuals naturally resistant to SCALA SC FUNGICIDE and other Group 9 fungicides. A gradual or total loss of disease control may occur over time if these fungicides are used repeatedly in the same fields. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as

enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay fungicide resistance:

- Where possible, rotate the use of SCALA SC FUNGICIDE or other Group 9 fungicides with different groups that control the same pathogens. Use tank mixtures with fungicides from a different group when such use is permitted.
- For *Botrytis* control, follow the Fungicide Resistance Action Committee (FRAC) guidelines:
  - Always alternate applications of SCALA SC FUNGICIDE or other Group 9 fungicides with fungicides having a different mode of action (different Group).
  - In situations where 3 to 6 *Botrytis* treatments are made per crop and season, a maximum of 2 applications of SCALA SC FUNGICIDE or any other Group 9 fungicide are recommended. In situations where 7 or more *Botrytis* treatments are made per crop and season, a maximum of 3 applications of SCALA SC FUNGICIDE or any other Group 9 fungicide are recommended.
- Avoid consecutive applications of SCALA SC FUNGICIDE or other Group 9 fungicides in a season. Alternate applications with fungicides having a different mode of action.
- Fungicide use should be based on an IPM program that includes scouting, historical information related to pesticide use and crop rotation and considers cultural, biological and other chemical control practices.
- Monitor treated fungal populations for resistance development. If disease continues to progress after treatment with this product, do not increase the use rate. Discontinue use of this product, and switch to another fungicide with a different target site of action, if available.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.
- For further information and to report suspected resistance, contact a Bayer CropScience representative at 1-888-283-6847 or at [www.bayercropscience.com](http://www.bayercropscience.com).

## **Section 10: Mixing Instructions**

### **MIXING INSTRUCTIONS:**

SCALA SC FUNGICIDE must be applied with properly calibrated, clean equipment.

Prior to adding SCALA SC FUNGICIDE to the spray tank, ensure that the spray tank is thoroughly clean.

- Add one-half of the required amount of water to the spray or mixing tank and start agitation.
- Add the required quantity of SCALA SC FUNGICIDE to the water and complete filling with water to the required total volume.
- Maintain agitation throughout mixing and spraying.
- When tank-mixing with BRAVO 500 for use on potatoes, follow use directions according to the BRAVO 500 Label.

## Section 11: Preharvest Intervals (PHI)

- Pre-harvest Intervals:
  - Apple (early season application) 72 day
  - Apple (late season application) 14 day
  - Grape 7 day
  - Pear 72 day
  - Strawberry 1 day
  - Potato 7 day
  - Greenhouse tomato 1 day

## Section 12: Notices

### NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

FOR MORE INFORMATION CONTACT:

Notification Change

Bayer CropScience Inc.  
~~#100-3131-114<sup>th</sup> Avenue SE~~  
Calgary, AB ~~T2Z 3X2~~

Suite 200, 160 Quarry Park Blvd. S.E.  
Calgary, Alberta T2C 3G3

SCALA is a registered trademark of Bayer.  
All other products mentioned are trademarks of their respective companies.

091116

\*\*\*\*\*

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.