

GROUP	9	12	FUNGICIDES
-------	---	----	------------

SWITCH™ 62.5 WG Fungicide

FUNGICIDE

AGRICULTURAL

Fungicide for control or suppression of listed diseases on onions, strawberries, blueberries (highbush and lowbush), raspberries, blackberries, carrots, mustard greens, turnip greens, cabbage, Saskatoon berries and dry and succulent beans.

GUARANTEE:

Cyprodinil 37.5%
Fludioxonil 25.0%

**READ THE LABEL AND BOOKLET BEFORE USING
KEEP OUT OF THE REACH OF CHILDREN**

CAUTION – EYE IRRITANT

REGISTRATION NO.: **28189**
PEST CONTROL PRODUCTS ACT

NET CONTENTS: **794 g - 1kg**

Syngenta Crop Protection Canada, Inc.
140 Research Lane, Research Park
Guelph, Ontario
N1G 4Z3
Telephone: 1-877-964-3682

Label

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

FIRST AID

IN CASE OF POISONING, call a physician or poison control centre **IMMEDIATELY**. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

If swallowed, call a poison control centre or doctor **IMMEDIATELY** for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing, take off contaminated clothing. Rinse skin **IMMEDIATELY** with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes, hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

TOXICOLOGICAL INFORMATION

There is no specific antidote known. Treat symptomatically. If large amount has been ingested and emesis is inadequate, lavage stomach.

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN. Avoid contact with eyes or prolonged contact with skin or clothing. Avoid inhalation of dust. Causes moderate eye irritation. Wear a long-sleeved shirt, long pants, shoes plus socks, coveralls and chemical resistant gloves such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, or Viton during mixing and loading. Wear a long-sleeved shirt, long pants and shoes plus socks during application, clean-up and repair. Wear goggles during application. **Do not re-enter treated area within 12 hours after application.** Do not eat, drink or smoke while mixing or during application. Change and wash clothing immediately after use. Wash hands and face before eating, drinking, smoking or using the toilet. Store and wash all protective clothing separately from household laundry. Wash clothing in detergent and hot water before reuse.

ENVIRONMENTAL PRECAUTIONS

This product is TOXIC to aquatic organisms. Observe buffer zones specified under GENERAL USE DIRECTIONS.

DO NOT apply this product directly to aquatic habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs, ditches, and wetlands), estuaries, or marine habitats.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

This product is persistent and will carryover. It is recommended that any products containing fludioxonil not be used in areas treated with this product during the previous season.

USE ONLY ON CROPS AS LISTED ON THIS LABEL.

DO NOT APPLY BY AIR.

Do not apply this product through any type of irrigation system.

STORE IN A DRY PLACE.

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's website at www.croplife.ca.

DECONTAMINATION AND DISPOSAL

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean up of spills.

CONTAINER DISPOSAL:

FOR DISPOSAL OF PLASTIC JUGS:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

***IN CASE OF EMERGENCY INVOLVING A MAJOR SPILL, FIRE OR POISONING,
CALL 1-800-327-8633 (FASTMED)***

GROUP	9	12	FUNGICIDES
-------	---	----	------------

SWITCH™ 62.5 WG Fungicide

FUNGICIDE

AGRICULTURAL

Fungicide for control or suppression of listed diseases on onions, strawberries, blueberries (highbush and lowbush), raspberries, blackberries, carrots, mustard greens, turnip greens, cabbage, Saskatoon berries and dry and succulent beans.

GUARANTEE:

Cyprodinil	37.5%
Fludioxonil	25.0%

**READ THE LABEL AND BOOKLET BEFORE USING
KEEP OUT OF THE REACH OF CHILDREN**

CAUTION – EYE IRRITANT

REGISTRATION NO.: **28189**
PEST CONTROL PRODUCTS ACT

Syngenta Crop Protection Canada, Inc.
140 Research Lane, Research Park
Guelph, Ontario
N1G 4Z3
Telephone: 1-877-964-3682

Pamphlet

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

FIRST AID

IN CASE OF POISONING, call a physician or poison control centre **IMMEDIATELY**. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

If swallowed, call a poison control centre or doctor **IMMEDIATELY** for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing, take off contaminated clothing. Rinse skin **IMMEDIATELY** with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes, hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

TOXICOLOGICAL INFORMATION

There is no specific antidote known. Treat symptomatically. If large amount has been ingested and emesis is inadequate, lavage stomach.

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN. Avoid contact with eyes or prolonged contact with skin or clothing. Avoid inhalation of dust. Causes moderate eye irritation. Wear a long-sleeved shirt, long pants, shoes plus socks, coveralls and chemical resistant gloves such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, or Viton during mixing and loading. Wear a long-sleeved shirt, long pants and shoes plus socks during application, clean-up and repair. Wear goggles during application. **Do not re-enter treated area within 12 hours after application.** Do not eat, drink or smoke while mixing or during application. Change and wash clothing immediately after use. Wash hands and face before eating, drinking, smoking or using the toilet. Store and wash all protective clothing separately from household laundry. Wash clothing in detergent and hot water before reuse.

ENVIRONMENTAL PRECAUTIONS

This product is TOXIC to aquatic organisms. Observe buffer zones specified under GENERAL USE DIRECTIONS.

DO NOT apply this product directly to aquatic habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs, ditches, and wetlands), estuaries, or marine habitats.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

This product is persistent and will carryover. It is recommended that any products containing fludioxonil not be used in areas treated with this product during the previous season.

USE ONLY ON CROPS AS LISTED ON THIS LABEL.

DO NOT APPLY BY AIR.

Do not apply this product through any type of irrigation system.

STORE IN A DRY PLACE.

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's website at www.croplife.ca.

DECONTAMINATION AND DISPOSAL

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean up of spills.

CONTAINER DISPOSAL:

FOR DISPOSAL OF PLASTIC JUGS:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

**IN CASE OF EMERGENCY INVOLVING A MAJOR SPILL, FIRE OR POISONING
CALL 1-800-327-8633 (FASTMED)**

GENERAL INFORMATION

SWITCH™ 62.5WG Fungicide is a broad-spectrum fungicide for the control of listed diseases.

FAILURE TO FOLLOW DIRECTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR DISEASE CONTROL, AND/OR ILLEGAL RESIDUES.

Spray Equipment

Thorough coverage is necessary to provide good disease control. Applications using sufficient water volume to provide thorough and uniform coverage generally provide the most effective disease control.

For ground application equipment, a minimum of 200 L of water per hectare is recommended.

To avoid spray drift, do not apply when conditions favor drift beyond the target area. Avoid spray overlap, as crop injury may occur.

Equip sprayers with nozzles that provide accurate and uniform application. Calibrate sprayer before use.

GENERAL USE DIRECTIONS

Field sprayer application: DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT apply with spray droplets smaller than the ASAE fine classification.

Buffer zones:

The buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs, and wetlands), and estuarine / marine habitats.

Method of application	Buffer zones (metres) required for the protection of aquatic habitats of depths:		
	Less than 1 m	1 - 3 m	Greater than 3 m
Field sprayer	2	1	0

Use a pump with capacity to maintain the correct rated pressure for the nozzles selected. Maintain sufficient agitation to keep the mixture in suspension. Use a jet agitator, liquid sparge tube, or mechanical paddle for agitation. Do not air sparge.

It is suggested that screens be used to prevent nozzles from clogging. Screens placed after the tank and before the nozzles should be 50-mesh or coarser. Check nozzle manufacturer's recommendations.

For more information on spray equipment and calibration, consult sprayer manufacturers and provincial recommendations.

Mixing Procedures

Prepare only the amount of spray mixture that is needed for the immediate operation. Thoroughly clean spray equipment before using this product. Vigorous agitation is necessary for proper dispersal of the product. Maintain maximum agitation throughout the spraying operation. Do not let the spray mixture stand overnight in the spray tank. Flush the spray equipment thoroughly following each use and apply the rinsate to a previously treated area.

1. Add 1/2 of the required amount of water to the mix tank.
2. With the agitator running, add the SWITCH 62.5WG Fungicide to the tank.
3. Continue agitation while adding the remainder of the water.
4. Begin application of the solution after the SWITCH 62.5WG Fungicide has completely dispersed into the mix water.
5. Maintain agitation until all of the mixture has been applied.

Rotational Crop Restrictions: Do not plant any other crop for a period of 12 months after harvest or crop failure unless SWITCH 62.5WG Fungicide is registered for that use.

NOTE TO BUYER / USER

The DIRECTIONS FOR USE for this product for the use described below were developed by persons other than Syngenta Crop Protection Canada, Inc. and accepted for registration by Health Canada under the User Requested Minor Use Registration program. Syngenta Crop Protection Canada, Inc. itself makes no representation or warranty with respect to performance (efficacy) and/or crop tolerance (phytotoxicity) claims for this product when used on the crops listed below.

Accordingly, the Buyer and User assume all liability arising, and agree to hold Syngenta Crop Protection Canada, Inc. harmless, from any claims based on efficacy and/or phytotoxicity in connection with the use described below.

DIRECTIONS FOR USE

Ground application only. DO NOT APPLY BY AIR.

Do not exceed 2.9 kg of product per hectare or 3 applications per year.

Apply the higher rate and shorter interval under conditions of high disease pressures.

Make no more than two (2) sequential applications of SWITCH 62.5 WG Fungicide before alternating with another registered fungicide with a different mode of action.

Crop	Diseases	Product Rate in g/ha	Application Timing
Onions: Dry bulb, green, and onions grown for seed	Control of Botrytis Leaf Blight or Blast (<i>Botrytis squamosa</i>) Suppression of Purple Blotch (<i>Alternaria porri</i>)	775 to 975	Begin applications when conditions become favourable for disease but before infection. If favourable conditions persist make additional applications on 7 to 14 day intervals. Do not apply within 7 days of harvest.
Strawberries	Control of Gray Mold (<i>Botrytis cinerea</i>)	975	Begin applications at or before bloom and continue on 7 to 10 day intervals. Pre-harvest interval is one (1) day.
Blueberries: (highbush and lowbush)	Control of Anthracnose (<i>Colletotrichum acutatum</i>), and Botrytis Fruit Rot (<i>Botrytis cinerea</i>)	775 to 975	Make the first application during early bloom. A second application may be made 7 to 10 days later. A third application can be made if conditions remain favourable for disease development. Pre-harvest interval is one (1) day.
Caneberries: Raspberries, Blackberries	Control of Botrytis Fruit Rot (<i>Botrytis cinerea</i>)	775 to 975	Make the first application during early bloom. A second application may be made 7 to 10 days later. A third application can be made if conditions remain favourable for disease development. Pre-harvest interval is one (1) day.
Carrots	Control of Alternaria Leaf Blight (<i>Alternaria dauci</i>)	775 to 975	The first application should be made when the disease first appears; a second application should be made 7-10 days later. One more application may be made a minimum of 7 days later if conditions remain favourable for disease development. Do not apply within 7 days of harvest. Apply in sufficient water volume to obtain thorough coverage; a minimum spray volume of 200 L/ha is recommended.
Mustard Greens, Turnip Greens	Control of Powdery Mildew (<i>Erysiphe polygoni</i> , <i>E. heraclei</i>)	775 to 975	The first application should be made when disease first appears and continue on 7 to 10 day intervals. Apply in sufficient water volume to obtain thorough coverage; a minimum spray volume of 200 L/ha is recommended. Do not apply within 7 days of harvest.
Cabbage	Suppression of Alternaria Leaf Blight (<i>Alternaria brassicicola</i>)	775 to 975	The first application should be made when disease first appears and continue on 7 to 10 day intervals. Apply in sufficient water volume to obtain thorough coverage; a minimum spray volume of 200 L/ha is recommended. Do not apply within 7 days of harvest.

Crop	Diseases	Product Rate in g/ha	Application Timing
Saskatoon berries	Suppression of Entomosporium leaf spot (<i>Entomosporium mespili</i>)	775 to 975	<p>The first application should be made during early bloom. A second application may be made 7 to 10 day later. A third application can be make if conditions remain favourable for disease development.</p> <p>Use higher rate and shorter application interval under conditions that promote rapid disease development or if there is a history of high disease pressure in the field.</p> <p>Apply in sufficient water to ensure thorough coverage (187-935 L/ha). Do not enter or allow worker entry into treated Saskatoon berry fields for hand-harvesting, pruning or thinning during the restricted entry interval (REI) of 10 days. For all other activities, do not enter or allow worker entry into treated areas within 12 hours after application. Do not apply within 1 day of harvest. Do not make more than 3 applications per year</p>
Beans (dried and succulent, except cowpea) Including: chickpea (garbanzo bean), beans (<i>Lupinus</i> spp. including grain lupin, sweet lupin, white lupin, white sweet lupin), beans (<i>Phaseolus</i> spp. including kidney bean, lima bean, mung bean, navy bean, pinto bean, snap bean, wax bean), broad bean (fava bean), beans (<i>Vigna</i> spp. asparagus bean, black-eyed pea)	Control of White mold (<i>Sclerotinia sclerotiorum</i>) and Grey mold (<i>Botrytis cinerea</i>)	775 - 975	<p>Begin applications prior to or at the onset of disease and repeat applications at 7 day intervals if conditions remain favourable for disease development.</p> <p>For white mold control, make the first application at 10-20% bloom. In some locations, a single application at this timing will provide adequate disease control.</p> <p>Apply the higher application rate under conditions of high disease pressure.</p> <p>Apply in sufficient water to ensure thorough coverage: Ground: 175 – 225 L of water /ha</p> <p>Do not apply more than 1.1 kg ai/ha of cyprodinil products and 0.73 kg ai/ha of fludioxonil containing products per crop.</p> <p>DO NOT enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.</p> <p>Do not apply within 7 days of harvest.</p>

Resistance Management Recommendations

For resistance management, SWITCH 62.5WG Fungicide contains a Group 9 and Group 12 Fungicide. Any fungal population may contain individuals naturally resistant to SWITCH 62.5WG Fungicide and other Group 9 and Group 12 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Other resistance mechanisms that are not linked to site of action but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance management strategies should be followed.

To delay fungicide resistance:

Where possible, rotate the use of SWITCH 62.5 WG Fungicide or other Group 9 and Group 12 fungicides with different groups that control the same pathogens.

Avoid application of more than the maximum number listed in the label and do not apply more than 2 consecutive sprays of SWITCH 62.5WG Fungicide or other fungicides in the same group in a season.

Fungicide use should be based on an IPM program that includes scouting, historical information related to pesticide use and crop rotation and considers cultural, biological and other chemical control practices.

Monitor treated fungal populations for sign of resistance development. If disease continues to progress after treatment with this product, do not increase the use rate. Discontinue use of this product, and switch to another fungicide with a different target site of action, if available.

Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for specific crops and disease problems in your area.

For further information or to report suspected resistance, contact company representatives at 1-87-SYNGENTA (1-877-964-3682) or at www.syngenta.ca.

SWITCH™ is a trademark of a Syngenta Group Company.

GROUP	9	12	FUNGICIDES
--------------	----------	-----------	-------------------

EMERGENCY USE LABEL

SWITCH® 62.5 WG Fungicide

FOR EMERGENCY USE ONLY

FUNGICIDE

AGRICULTURAL

For sale and use in Ontario to control powdery mildew on greenhouse cucumbers until December 31, 2009.

COMMERCIAL

GUARANTEE:

Cyprodinil	37.5%
Fludioxonil	25.0%

**READ THE LABEL AND BOOKLET BEFORE USING
KEEP OUT OF THE REACH OF CHILDREN**

CAUTION – EYE IRRITANT

REGISTRATION NO.: **28189**
PEST CONTROL PRODUCTS ACT

NET CONTENTS: **794 g – 1 kg**

Syngenta Crop Protection Canada, Inc.
140 Research Lane, Research Park
Guelph, Ontario
N1G 4Z3 Telephone: 1-877-964-3682

GENERAL INFORMATION

SWITCH 62.5WG Fungicide is a broad-spectrum fungicide for the control of powdery mildew in greenhouse cucumbers.

Chemigation

Do not apply this product through any type of irrigation system.

FAILURE TO FOLLOW DIRECTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR DISEASE CONTROL, AND/OR ILLEGAL RESIDUES.**NOTE TO USER:**

The DIRECTIONS FOR USE for this product for the use described in this emergency use label were developed by persons other than Syngenta Crop Protection Canada, Inc. and accepted for registration by Health Canada under the Emergency Registration. Syngenta Crop Protection Canada, Inc. itself makes no representation or warranty with respect to performance (efficacy) and/or crop tolerance (phytotoxicity) claims for this product when used on the crops recommended below.

Accordingly, the User assumes all liability arising, and agrees to hold Syngenta Crop Protection Canada, Inc. harmless, from any claims based on efficacy and/or phytotoxicity in connection with the use described in this label.

PROPOSED DIRECTIONS FOR NEW USE

Product Name / Reg. No.	SWITCH 62.5 WG Fungicide / Reg. No. 28189
Pest(s)	Powdery mildew (<i>Podosphaera xanthii</i>).
Site (Crop)	Cucumbers grown in the greenhouse.
Method of Application	Ground application only.
Rate of Application and Spray Volume (include adjuvant rate, if being proposed)	775 g product/ha. Applied in sufficient water to ensure thorough coverage (200 - 3000 L/ha).
Number of Applications	Do not make more than two (2) applications per year
Application Interval	Alternate with other registered fungicides on 7 to 10 days schedule.
Application Timing	The first application should be made when disease first appears; a second application should be made 7-10 days later.
Pre-Harvest Interval	Do not apply within 2 days of harvest.
Other	Make no more than two (2) sequential applications of SWITCH 62.5 WG Fungicide before alternating with another registered fungicide with a different mode of action.

FIRST AID

IN CASE OF POISONING, call a physician or poison control centre **IMMEDIATELY**. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

IF SWALLOWED, call a poison control centre or doctor **IMMEDIATELY** for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF ON SKIN/CLOTHING, take off contaminated clothing. Rinse skin **IMMEDIATELY** with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

IF IN EYES, hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

IF INHALED, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.

TOXICOLOGICAL INFORMATION

There is no specific antidote known. Treat symptomatically. If large amount has been ingested and emesis is inadequate, lavage stomach.

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN. Avoid contact with eyes or prolonged contact with skin or clothing. Avoid inhalation of dust. Causes moderate eye irritation. Wear a long-sleeved shirt, long pants and gloves during mixing, loading, application, clean-up and repair. Wear goggles during application. When using a backpack sprayer, wear gloves during application. **Do not re-enter treated area within 12 hours after application**. Do not eat, drink or smoke while mixing or during application. Change and wash clothing immediately after use. Wash hands and face before eating, drinking, smoking or using the toilet. Store and wash all protective clothing separately from household laundry. Wash clothing in detergent and hot water before reuse.

ENVIRONMENTAL PRECAUTIONS

This product is TOXIC to aquatic organisms. Observe buffer zones specified under GENERAL USE DIRECTIONS.

DO NOT apply this product directly to aquatic habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs, ditches, and wetlands), estuaries, or marine habitats.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

This product is persistent and will carryover. It is recommended that any products containing fludioxonil not be used in areas treated with this product during the previous season.

DO NOT allow effluent or runoff from greenhouses or mushroom houses containing this product to enter lakes, streams, ponds or other waters.

GENERAL USE DIRECTIONS

Spray Equipment

Thorough coverage is necessary to provide good disease control. Applications using sufficient water volume to provide thorough and uniform coverage generally provide the most effective disease control.

For ground application equipment, a minimum of 200 L of water per hectare is recommended.

To avoid spray drift, do not apply when conditions favor drift beyond the target area. Avoid spray overlap, as crop injury may occur.

Equip sprayers with nozzles that provide accurate and uniform application. Calibrate sprayer before use.

Field sprayer application: DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT apply with spray droplets smaller than the ASAE fine classification.

Buffer zones:

The buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs, and wetlands), and estuarine / marine habitats.

Method of application	Buffer zones (metres) required for the protection of aquatic habitats of depths:		
	Less than 1 m	1 - 3 m	Greater than 3 m
Field sprayer	2	1	0

Mixing Procedures

Prepare only the amount of spray mixture that is needed for the immediate operation. Thoroughly clean spray equipment before using this product. Vigorous agitation is necessary for proper dispersal of the product. Maintain maximum agitation throughout the spraying operation. Do not let the spray mixture stand overnight in the spray tank. Flush the spray equipment thoroughly following each use and apply the rinsate to a previously treated area.

1. Add 1/2 of the required amount of water to the mix tank.
2. With the agitator running, add the SWITCH 62.5WG Fungicide to the tank.
3. Continue agitation while adding the remainder of the water.

4. Begin application of the solution after the SWITCH 62.5WG Fungicide has completely dispersed into the mix water.
5. Maintain agitation until all of the mixture has been applied.

Use a pump with capacity to maintain the correct rated pressure for the nozzles selected. Maintain sufficient agitation to keep the mixture in suspension. Use a jet agitator, liquid sparge tube, or mechanical paddle for agitation. Do not air sparge.

It is suggested that screens be used to prevent nozzles from clogging. Screens placed after the tank and before the nozzles should be 50-mesh or coarser. Check nozzle manufacturer's recommendations.

For more information on spray equipment and calibration, consult sprayer manufacturers and provincial recommendations.

Resistance Management Recommendations

For resistance management, SWITCH 62.5WG Fungicide contains a Group 9 and Group 12 Fungicide. Any fungal population may contain individuals naturally resistant to SWITCH 62.5WG Fungicide and other Group 9 and Group 12 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Other resistance mechanisms that are not linked to site of action but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance management strategies should be followed.

To delay fungicide resistance:

Where possible, rotate the use of SWITCH 62.5WG Fungicide or other Group 9 and Group 12 fungicides with different groups that control the same pathogens.

Avoid application of more than the maximum number listed in the label and do not apply more than 2 consecutive sprays of SWITCH 62.5WG Fungicide or other fungicides in the same group in a season.

Fungicide use should be based on an IPM program that includes scouting, historical information related to pesticide use and crop rotation and considers cultural, biological and other chemical control practices.

Monitor treated fungal populations for sign of resistance development. If disease continues to progress after treatment with this product, do not increase the use rate. Discontinue use of this product, and switch to another fungicide with a different target site of action, if available.

Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for specific crops and disease problems in your area.

For further information or to report suspected resistance, contact company representatives at 1-877-SYNGENTA (1-877-964-3682) or at www.syngenta.ca

STORE IN A DRY PLACE.

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's web site at www.croplife.ca.

DECONTAMINATION AND DISPOSAL

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean up of spills.

CONTAINER DISPOSAL:

FOR DISPOSAL OF PLASTIC JUGS:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

***IN CASE OF EMERGENCY INVOLVING A MAJOR SPILL, FIRE OR POISONING,
CALL 1-800-327-8633 (FASTMED)***

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

Product names marked ® or ™ are trademarks of a Syngenta Group Company

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.