

ACCORD WSB HERBICIDE

GUARANTEE: Quinclorac ... 75%

Warning, contains the allergen sulfites

REGISTRATION NO. 27813  
PEST CONTROL PRODUCTS ACT

NET CONTENTS: 135 g - 270 g



CAUTION

POISON

CAUTION - EYE AND SKIN IRRITANT  
POTENTIAL SKIN SENSITIZER

READ THE LABEL AND ATTACHED PAMPHLET BEFORE USING.  
KEEP OUT OF REACH OF CHILDREN.  
DO NOT SELL WATER-SOLUBLE BAGS SEPARATELY.

**ACCORD® Water-Soluble Bag (WSB) Herbicide**

For selective post-emergence control of green foxtail, cleavers, volunteer flax, barnyard grass and suppression of annual and perennial sow-thistle in spring and durum wheat, spring barley (grown for feed purposes only) and canary seed.

FOR SALE FOR USE IN THE PRAIRIE PROVINCES AND PEACE RIVER REGION OF BRITISH COLUMBIA ONLY

**COMMERCIAL (AGRICULTURAL)**

**GUARANTEE:** Quinclorac ..... 75%

**Warning, contains the allergen sulfites**

**REGISTRATION NO.** 27813

**PEST CONTROL PRODUCTS ACT**

**IN CASE OF EMERGENCY ENDANGERING LIFE OR PROPERTY  
INVOLVING THIS PRODUCT, CALL DAY OR NIGHT  
1-800-454-2673**



**CAUTION**

**POISON**

**CAUTION - EYE AND SKIN IRRITANT  
POTENTIAL SKIN SENSITIZER**

**NET CONTENTS:** 0.54 - 1.08 kg (4 x 135 - 270 grams water-soluble bags)

**READ THE LABEL AND ATTACHED PAMPHLET BEFORE USING.  
KEEP OUT OF REACH OF CHILDREN.  
DO NOT SELL WATER-SOLUBLE BAGS SEPARATELY.**

BASF Canada Inc.  
100 Milverton Drive  
5<sup>th</sup> Floor  
Mississauga, Ontario  
L5R 4H1  
1-877-371-2273

EPA Est. No.: 65387-AR-001

**ACCORD®** is a registered trademark of BASF Canada Inc.

## PRECAUTIONS

1. **KEEP OUT OF REACH OF CHILDREN.**
2. Do not take internally.
3. Avoid inhalation of vapour, dust or spray mist and contact with eyes, skin or clothing.
4. Wash thoroughly after handling and before eating, drinking or smoking.
5. Wear protective equipment and clothing, including goggles or face shield, gloves (rubber, PVC, neoprene or nitrile), hat, long-sleeved shirt, trousers and rubber boots during mixing, loading, application, clean-up and repair activities.
6. If clothing becomes contaminated, remove and wash separately from household laundry before reuse.
7. Clean spray equipment thoroughly after use. Refer to sprayer clean-up section.
8. Do not contaminate domestic or irrigation water, lakes, streams or ponds by the cleaning of equipment or the disposal of wastes.

The enclosed pouches of **ACCORD WSB herbicide** will dissolve in water. DO NOT allow the pouches to become wet before use. DO NOT handle the individual pouches with wet hands or wet gloves. Always reseal the overwrap bag to protect the remaining unused pouches. The entire contents of the inner water soluble bag must be used.

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's web site at [www.croplife.ca](http://www.croplife.ca).

## FIRST AID

**If swallowed:** Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

**If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

**If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

## TOXICOLOGICAL INFORMATION

Treat symptomatically.

## STORAGE

1. Store in original, tightly-closed container.
2. Do not ship or store near food, feed, seed or fertilizers.
3. Store in cool, dry, locked, well-ventilated area without floor drain.
4. Herbicides should be shipped or stored separately from other pesticides to avoid cross-contamination.
5. Freezing will not harm **ACCORD WSB herbicide**. Should product freeze, warm to room temperature prior to use.

## DISPOSAL

1. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
2. Make the empty container unsuitable for further use.
3. Dispose of the container in accordance with provincial requirements.
4. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

## NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

**ACCORD® Water-Soluble Bag (WSB) Herbicide**

For selective post-emergence control of green foxtail, cleavers, volunteer flax, barnyard grass and suppression of annual and perennial sow-thistle in spring and durum wheat, spring barley (grown for feed purposes only) and canary seed.

FOR SALE FOR USE IN THE PRAIRIE PROVINCES AND PEACE RIVER REGION OF BRITISH COLUMBIA ONLY

**COMMERCIAL (AGRICULTURAL)**

**GUARANTEE:** Quinclorac ..... 75%

Warning, contains the allergen sulfites

**REGISTRATION NO.** 27813

**PEST CONTROL PRODUCTS ACT**

**IN CASE OF EMERGENCY ENDANGERING LIFE OR PROPERTY  
INVOLVING THIS PRODUCT, CALL DAY OR NIGHT  
1-800-454-2673**



**CAUTION**

**POISON**

**CAUTION - EYE AND SKIN IRRITANT  
POTENTIAL SKIN SENSITIZER**

**NET CONTENTS: 0.54 - 1.08 kg (4 x 135 - 270 grams water-soluble bags)**

**READ THE LABEL AND ATTACHED PAMPHLET BEFORE USING.  
KEEP OUT OF REACH OF CHILDREN.  
DO NOT SELL WATER-SOLUBLE BAGS SEPARATELY.**

BASF Canada Inc.  
100 Milverton Drive  
5<sup>th</sup> Floor  
Mississauga, Ontario  
L5R 4H1  
1-877-371-2273

EPA Est. No.: 65387-AR-001

**ACCORD®** is a registered trademark of BASF Canada Inc.

## GENERAL INFORMATION

**ACCORD WSB herbicide** is a herbicide for selective post-emergence control of green foxtail (including Group 1 and Group 3 resistant biotypes), volunteer flax, cleavers, barnyard grass and suppression of annual and perennial sow-thistle in hard red spring, Canadian prairie spring, durum, Canada Western extra strong wheats, spring barley (grown for feed purposes only) and canary seed.

**ACCORD WSB herbicide** is a herbicide with mainly systemic action. Uptake into the plant occurs through both the foliage and root system. Thorough coverage of foliage is important for consistent weed control. Failure to penetrate crop or weed leaf canopies with the spray will result in inconsistent control of weeds growing underneath.

Visual symptoms of weed control of **ACCORD WSB herbicide** may take up to two weeks following application to develop. These symptoms include initial twisting to stunting, reddening and chlorosis about 14 days followed by necrosis and death about 21 days after application. Even though **ACCORD WSB herbicide** symptoms may take some time to develop, competition from the weeds treated with **ACCORD WSB herbicide** is eliminated soon after application.

## DIRECTIONS FOR USE

### Application Rate and Timing for Wheat, Spring Barley (grown for feed purposes only) and Canary Seed

DO NOT APPLY BY AIR.

Apply **ACCORD WSB herbicide** at 135-165 g/ha when weeds are small and actively growing. **ACCORD WSB herbicide** will control the following weeds at the timing detailed in **Table 1**. **ACCORD WSB herbicide** can be applied to wheat, spring barley and canary seed at the maximum application rates and timing detailed in **Table 1**.

Use the 135 g/ha rate ONLY for control of volunteer flax, barnyard grass, cleavers, lighter infestations of green foxtail and suppression of annual and perennial sow-thistle. Use the higher rate of 165 g/ha for control of heavier infestations of green foxtail. (Do not use the 165 g/ha rate on barley.) **Use only the 135 g/ha rate when applying ACCORD WSB herbicide to spring barley.**

DO NOT apply **ACCORD WSB herbicide** to any field more often than every second year. This practice must be respected in order to avoid potential injury to future rotational crops, to minimize the potential for carryover and accumulation of soil residues, and to reduce the selection pressure which could contribute to the development of resistant biotypes.

Early treatment of weeds with **ACCORD WSB herbicide** is important to maximize crop yield potential through elimination of early weed competition. Some initial crop injury may be observed after application, but this is usually outgrown and should not affect crop yield.

**TABLE 1: WEED APPLICATION TIMING TABLE**

<b>WEED</b>	<b>RANGE</b>
Green foxtail	1 - 5 leaf max 2 tillers
Volunteer flax	1 - 8 cm.
Cleavers	1 - 3 whorls
Barnyard grass	1 - 5 leaf
Annual sow-thistle*	2 - 6 leaf
Perennial sow-thistle*	2 - 6 leaf
<b>CROP</b>	
Spring barley** (grown for feed purposes only)	1 - 4 leaf (prior to tillering)
Wheat (spring and durum)	1 - 5 leaf
Canary seed***	3 - 5 leaf

\* Suppression only.

\*\* Maximum rate for barley is 135 g/ha. To avoid crop injury apply **ACCORD WSB herbicide** before spring barley tillers.

\*\*\* Not to be used for human consumption or fed to livestock.

**All leaf stages listed above refer to true leaves.**

## **SPRAYING INSTRUCTIONS**

### **Ground Application**

Use sprayers equipped with standard flat fan pesticide nozzles with a spray volume of 100 L/ha at a constant pressure of 275-425 kPa. Tilt spray nozzles 45 degrees forward to ensure better coverage. The use of 50 mesh strainers and screens is recommended.

### **ADDITIVES**

Always use **MERGE** adjuvant at 1.0% v/v for optimum performance of **ACCORD WSB herbicide**.

**TANK MIX COMBINATIONS  
(BARLEY AND WHEAT APPLICATIONS ONLY):**

For the appropriate rate of **ACCORD WSB herbicide**, refer to the Application Rate and Timing section of the label.

**Broadleaf Weed Control**

Although **ACCORD WSB herbicide** provides control of several broadleaf weeds, a tank mix with a broadleaf compound is required to give broad spectrum broadleaf weed control.

For control of additional broadleaf weeds in barley and wheat, **ACCORD WSB herbicide** can be tank mixed with any of the broadleaf herbicides listed in **Table 2**. When tank mixing **ACCORD WSB herbicide** with these broadleaf herbicides, a slight reduction in control of green foxtail may be observed. The level of green foxtail control may be improved by using the 165 g/ha rate of **ACCORD WSB herbicide** in the tank mixture.

Refer to **Table 2** for appropriate use rates and timing of crop applications. Always refer to the broadleaf herbicide label for additional precautions, restrictions and use instructions.

**TABLE 2: TANK MIX OPTIONS FOR ACCORD WSB HERBICIDE  
IN BARLEY AND WHEAT**

Crop	ACCORD WSB Herbicide Rate (g/L)	Tank Mix		Crop Stage	Weeds Controlled
		Partner	Rate		
Barley	135 - 165	<b>MCPA Amine</b> (assume 500 series)	1.1 L/ha	3-4 leaf	<b>ACCORD WSB Herbicide</b> - see Table 1. Broadleaf weeds listed on <b>MCPA Amine</b> label.
		<b>MCPA Ester</b> (assume 500 series)	1.1 L/ha	3-4 leaf	<b>ACCORD WSB Herbicide</b> - see Table 1. Broadleaf weeds listed on <b>MCPA Ester</b> label.
		<b>Buctril M</b>	1.0 L/ha	2-4 leaf	<b>ACCORD WSB Herbicide</b> - see Table 1. Broadleaf weeds listed on <b>Buctril M</b> label.
		<b>Refine Extra<sup>1</sup></b>	20 g/ha	2-4 leaf	<b>ACCORD WSB Herbicide</b> - see Table 1. Broadleaf weeds listed on <b>Refine Extra</b> label.
Wheat (spring and durum)	135 - 165	<b>Buctril M</b>	1.0 L/ha	2-5 leaf	<b>ACCORD WSB Herbicide</b> - see Table 1. Broadleaf weeds listed on <b>Buctril M</b> label.
		<b>2,4-D Amine</b> (assume 500 series)	0.840 - 1.1 L/ha	3-5 leaf	<b>ACCORD WSB Herbicide</b> - see Table 1. Broadleaf weeds listed on <b>2,4-D Amine</b> label.
		<b>2,4-D Ester</b> (assume 500 series)	0.840 - 1.1 L/ha	4-5 leaf	<b>ACCORD WSB Herbicide</b> - see Table 1. Broadleaf weeds listed on <b>2,4-D Ester</b> label.
		<b>MCPA Amine</b> (assume 500 series)	0.840 - 1.1 L/ha	3-5 leaf	<b>ACCORD WSB Herbicide</b> - see Table 1. Broadleaf weeds listed on <b>MCPA Amine</b> label.
		<b>MCPA Ester</b> (assume 500 series)	0.840 - 1.1 L/ha	3-5 leaf	<b>ACCORD WSB Herbicide</b> - see Table 1. Broadleaf weeds listed on <b>MCPA Ester</b> label.
		<b>Refine Extra<sup>1</sup></b>	20 g/ha	2-5 leaf	<b>ACCORD WSB Herbicide</b> - see Table 1. Broadleaf weeds listed on <b>Refine Extra</b> label.
		<b>Express Pack<sup>1</sup></b> (Express + 2,4-D)	10 g/ha + 0.625 L/ha	3-5 leaf	<b>ACCORD WSB Herbicide</b> - see Table 1. Broadleaf weeds listed on <b>Express Pack</b> label.

Crop	ACCORD WSB Herbicide Rate (g/L)	Tank Mix		Crop Stage	Weeds Controlled
		Partner	Rate		
Wheat (spring only)	135 - 165	<b>Avenge 200-C</b>	3.5 L/ha	1-5 leaf	<b>ACCORD WSB Herbicide</b> - see Table 1. Wild oats as listed on <b>Avenge 200-C</b> label.
		<b>Avenge 200-C + Buctril M</b>	3.5 L/ha + 1.0 L/ha	2-5 leaf	<b>ACCORD WSB Herbicide</b> - see Table 1. Broadleaf weeds listed on <b>Buctril M</b> label. Wild oats as listed on <b>Avenge 200-C</b> label.
		<b>Avenge 200-C + 2,4-D Ester</b> (assume 500 series)	3.5 L/ha + 0.840 - 1.1 L/ha	4-5 leaf	<b>ACCORD WSB Herbicide</b> - see Table 1. Broadleaf weeds listed on <b>2,4-D Ester</b> label. Wild oats as listed on <b>Avenge 200-C</b> label.
		<b>Avenge 200-C + MCPA Ester</b> (assume 500 series)	3.5 L/ha + 0.840 - 1.1 L/ha	3-5 leaf	<b>ACCORD WSB Herbicide</b> - see Table 1. Broadleaf weeds listed on <b>MCPA Ester</b> label. Wild oats as listed on <b>Avenge 200-C</b> label.
		<b>Avenge 200-C + Refine Extra<sup>1</sup></b>	3.5 L/ha + 20 g/ha	2-5 leaf	<b>ACCORD WSB Herbicide</b> - see Table 1. Broadleaf weeds listed on <b>Refine Extra</b> label. Wild oats as listed on <b>Avenge 200-C</b> label.

<sup>1</sup> Addition of surfactants other than **MERGE** adjuvant is not required.

Always refer to the tank mix partner label(s) for additional precautions, restrictions, use instructions and recropping information.

Do not delay spraying broadleaf weeds if grassy weeds are not in the correct stage for treatment. If green foxtail, wild oats and broadleaf weeds are not in the correct stages for treatment, apply separate applications of each herbicide timed to control the required spectrum of weeds. Use **MERGE** adjuvant only with all tank mixtures.

### Wild Oat Control

For additional control of wild oats when populations are between 1-200 plants per square metre, **ACCORD WSB herbicide** can be tank mixed with **AVENGE® 200-C** herbicide at the rate of 3.5 L/ha. USE ONLY ON SPRING WHEAT VARIETIES LISTED ON THE **AVENGE 200-C** HERBICIDE LABEL.

**AVENGE 200-C** herbicide can cause some crop injury. Refer to the **Table 3** for appropriate use rates and timing of crop application. Always refer to the **AVENGE 200-C** herbicide label for additional precautions, restrictions and use instructions.

**TABLE 3: TANK MIXING FOR WILD OAT CONTROL**

<b>Tank Mix Partner</b>	<b>Product Rate</b>	<b>Crop Stage</b>
<b>Avenge 200-C</b> herbicide	3.5 L/ha	1-5 leaf stage

#### **Wild Oats and Broadleaf Weed Control**

For control of wild oats when populations are between 1-200 plants per square metre and certain broadleaf weeds, **ACCORD WSB herbicide** can be tank mixed with **AVENGE 200-C** herbicide at the rate of 3.5 L/ha and one of the broadleaf herbicides listed in Table 2. USE ONLY ON SPRING WHEAT VARIETIES LISTED ON THE **AVENGE 200-C** HERBICIDE LABEL.

**AVENGE 200-C** herbicide can cause some crop injury. Refer to **Table 3** for appropriate use rates and timing of crop application. Always refer to both the **AVENGE 200-C** herbicide and the broadleaf herbicide label for additional precautions, restrictions and use instructions.

#### **Wheat Tank Mixing**

Although **ACCORD WSB herbicide** provides control of several broadleaf weeds, a tank mix with a broadleaf compound is required to give broad spectrum broadleaf weed control in wheat.

When tank mixing **ACCORD WSB herbicide** with these broadleaf herbicides, a slight reduction in control of green foxtail may be observed. The level of green foxtail control may be improved by using the 165 g/ha rate of **ACCORD WSB herbicide** in the tank mixture.

For additional control of wild oats when populations are between 1-200 plants per square metre, **ACCORD WSB herbicide** can be tank mixed with **AVENGE® 200-C** at the rate of 3.5 L/ha. USE ONLY ON SPRING WHEAT VARIETIES LISTED ON THE **AVENGE 200-C** LABEL. **AVENGE 200-C** can cause some crop injury.

#### **RECRIPPING**

Due to the residual activity of **ACCORD WSB herbicide** in the soil, land treated with **ACCORD WSB herbicide** cannot be rotated to crops other than specified in **Table 4**. To avoid injury to rotational crops, the minimum recropping intervals in **Table 4** must be followed.

**TABLE 4: MINIMUM RECROPPING INTERVALS**

<b>CROP</b>	<b>MINIMUM RECROPPING INTERVAL (Months)</b>	<b>NOTES</b>
Wheat* (spring, durum) Spring barley*	0 0	These crops can be re-planted in the same season as <b>ACCORD WSB herbicide</b> applications.
Canola Field peas Sunflowers	10 10 10	These crops and the crops listed above can be planted the year following application of <b>ACCORD WSB herbicide</b> .
<b>Oats</b>	<b>12</b>	<b>These crops and the crops listed above can be planted the year following application of <b>ACCORD WSB herbicide</b>.</b>
Flax Lentils	22 22	These crops and the crops listed above can be planted two years following application of <b>ACCORD WSB herbicide</b> .

\*In the event of crop failure, only spring or durum wheat or spring barley may be reseeded in fields treated with **ACCORD WSB herbicide**.

**ACCORD WSB herbicide should not be used on land where potatoes or vegetables are part of the rotation.**

The company recommends that a field bioassay (a test strip grown to maturity) be conducted the year before growing any crops other than those listed in **Table 4**.

On lighter soils with low organic matter or under dry conditions, some crop injury may occur particularly in flax and lentils but will not reduce yield. Under these conditions, the minimum recropping interval for flax and lentils should be extended by 12 months.

**Refer to the broadleaf or AVENGE 200-C herbicide label for specific additional recropping restrictions.**

**RESTRICTIONS AND LIMITATIONS**

1. Do not apply **ACCORD WSB herbicide** when weather conditions may cause spray drift from treated areas to adjacent crops. Certain crops such as alfalfa, clover species, fababeans, flax, lentils, ornamentals, potatoes and vegetables will be injured by spray drift of **ACCORD WSB herbicide**.
2. Do not treat any crops other than wheat, spring barley and canary seed.
3. Do not apply **ACCORD WSB herbicide** to wheat under seeded to forages.

4. Do not apply **ACCORD WSB herbicide** to wheat, spring barley or canary seed that has been subjected to stress from conditions such as frost, hail damage, flooding, drought, extended cold period, etc.
5. Rainfall within 6 hours after application may reduce effectiveness of spray.
6. When **ACCORD WSB herbicide** is applied beyond the recommended growth stages, limited crop injury and/or unsatisfactory weed control may result.
7. Cool weather conditions or drought will delay herbicide activity and if prolonged, may result in poor weed control.
8. Do not use **ACCORD WSB herbicide** with additives, pesticides or fertilizers not specifically recommended on this label.
9. Allow 4 days between application of **ACCORD WSB herbicide** and any other chemical not recommended as a tank mix combination on this label.
10. **ACCORD WSB herbicide** must not be applied within 77 days of harvest of wheat and within 80 days of harvest of spring barley.
11. Canary seed treated with **ACCORD WSB herbicide** is NOT TO BE USED FOR HUMAN CONSUMPTION OR FED TO LIVESTOCK.
12. Apply using ground equipment only. DO NOT APPLY BY AIR.
13. Overspray or drift into important wildlife habitats such as shelterbelts, wetlands, woodlots, vegetated ditch, ponds and lake banks and other cover on the edges of fields should be avoided. A 10-metre buffer zone should be observed adjacent to aquatic habitats such as streams, ponds, rivers and lakes and to areas that drain into these habitats. When a tank mixture is used, consult the label of the tank mix partners and observe the largest (most restrictive) buffer zone of the products involved in the tank mixture.
14. Do not graze the treated crops or cut for hay within 77 days of application.

## **MIXING**

1. Thoroughly clean the sprayer prior to use. For appropriate cleaning instructions, refer to the label of the product sprayed previous to application of **ACCORD WSB herbicide**.
2. Fill the clean spray tank half full with clean water. Start agitation or by-pass system. Agitation should be running during the entire mixing procedure.
3. Add the correct amount of **ACCORD WSB herbicide** and agitate 2 to 3 minutes.

4. Add the correct amount of broadleaf herbicide, followed by **AVENGE 200-C** herbicide, if required. When mixing **ACCORD WSB herbicide** with **EXPRESS PACK**, **EXPRESS** herbicide must be completely in suspension in the spray tank prior to adding **2,4-D** herbicide.

**NOTE:** On repeat tank loads of either **EXPRESS PACK** or **REFINE EXTRA** herbicide, prepare an **EXPRESS/water** or **REFINE/water** slurry in a separate container with clean water before adding to the spray tank.

5. Add the correct amount of **MERGE** adjuvant and agitate 2 to 3 minutes.
6. Add remainder of water to the spray tank and maintain sufficient agitation during mixing and spraying to ensure a uniform spray mixture.
7. For sprayer clean-up, refer to the sprayer clean-up section.
8. Consult the broadleaf or **AVENGE 200-C** herbicide label for additional application instructions, use precautions and recropping information.

#### **SPRAYER CLEAN-UP**

Certain crops such as alfalfa, clover species, fababeans, flax, lentils, ornamentals, potatoes, tomatoes and vegetables are particularly sensitive to **ACCORD WSB herbicide**. To avoid injury to subsequent crops other than wheat, the sprayer should be thoroughly cleaned immediately after use and prior to spraying other crops by performing the following steps:

1. Following spray application, drain any remaining spray solution, then flush the tank, boom and hoses with clean water until any visible residues are removed. (Repeat step 1, if necessary.) **DO NOT CLEAN SPRAYER NEAR DESIRABLE VEGETATION OR NEAR WELL OR WATER SOURCE.**
2. Completely fill spray tank with clean water while adding 1 litre of household ammonia (containing 3% ammonia) per 100 litres of water or a commercially licensed tank cleaner such as **FINNISH®**. Reduce the amount of ammonia added proportionally if higher concentrations (%) of ammonia are used. Flush the solution through the boom and nozzles and then add more water to completely refill the tank. Agitate the solution for at least 15 minutes and then flush the boom and nozzles until the spray tank is empty.
3. Remove the nozzles and screens and clean separately in a bucket containing a cleaning agent and water.
4. Repeat step 2.
5. Thoroughly rinse the tank with clean water and flush the water through the boom.

## RESISTANCE-MANAGEMENT RECOMMENDATIONS

For resistance management, **ACCORD WSB herbicide** is a Group 4 herbicide. Any weed population may contain or develop plants naturally resistant to **ACCORD WSB herbicide** and other Group 4 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

- Where possible, rotate the use of **ACCORD WSB herbicide** or other Group 4 herbicides with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group when such use is permitted.
- Herbicide use should be based on an IPM program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical), cultural, biological and other chemical control practices.
- Monitor treated weed populations for resistance development.
- Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment and planting clean seed.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact BASF at 1-877-371-2273.

## PRECAUTIONS

1. **KEEP OUT OF REACH OF CHILDREN.**
2. Do not take internally.
3. Avoid inhalation of vapour, dust or spray mist and contact with eyes, skin or clothing.
4. Wash thoroughly after handling and before eating, drinking or smoking.
5. Wear protective equipment and clothing, including goggles or face shield, gloves (rubber, PVC, neoprene or nitrile), hat, long-sleeved shirt, trousers and rubber boots during mixing, loading, application, clean-up and repair activities.
6. If clothing becomes contaminated, remove and wash separately from household laundry before reuse.
7. Clean spray equipment thoroughly after use. Refer to sprayer clean-up section.

8. Do not contaminate domestic or irrigation water, lakes, streams or ponds by the cleaning of equipment or the disposal of wastes.

The enclosed pouches of **ACCORD WSB herbicide** will dissolve in water. DO NOT allow the pouches to become wet before use. DO NOT handle the individual pouches with wet hands or wet gloves. Always reseal the overwrap bag to protect the remaining unused pouches. The entire contents of the inner water soluble bag must be used.

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's web site at [www.croplife.ca](http://www.croplife.ca).

## FIRST AID

**If swallowed:** Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

**If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

**If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

## TOXICOLOGICAL INFORMATION

Treat symptomatically.

## STORAGE

1. Store in original, tightly-closed container.
2. Do not ship or store near food, feed, seed or fertilizers.
3. Store in cool, dry, locked, well-ventilated area without floor drain.
4. Herbicides should be shipped or stored separately from other pesticides to avoid cross-contamination.

5. Freezing will not harm **ACCORD WSB herbicide**. Should product freeze, warm to room temperature prior to use.

## **DISPOSAL**

1. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
2. Make the empty container unsuitable for further use.
3. Dispose of the container in accordance with provincial requirements.
4. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

## **NOTICE TO USER**

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

**MERGE®** is a registered trademark of BASF Canada Inc.

®All other products listed are registered trademarks of their respective companies.

\*\*\*\*\*

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.