

2009-1604
29-JUN-2009

Water-soluble bag label E

NIPPON SODA

TRISTAR 70 WSP Insecticide

GUARANTEE: Acetamiprid.....70% by weight



READ THE LABEL AND BOOKLET BEFORE USING

KEEP OUT OF REACH OF CHILDREN

NET CONTENTS: 16 g

Overpack label

To Open

Tear Here

This outer bag is not water soluble.
Do not put in Sprayer Tank.

NIPPON SODA

GROUP	4	INSECTICIDE
--------------	----------	--------------------

TRISTAR 70 WSP Insecticide

Contains Acetamiprid

AGRICULTURAL

GUARANTEE: Acetamiprid.....70% by weight



READ THE LABEL AND BOOKLET BEFORE USING

KEEP OUT OF REACH OF CHILDREN

REGISTRATION NO. 27127 PEST CONTROL PRODUCTS ACT

Registered by:

Nippon Soda Co., Ltd.
2-1, 2-Chome Ohtemachi
Chiyoda-ku, Tokyo 100-8165,
Japan

Agent: David Latter
D. W. Latter and Associates
44 Tamarack Circle
Toronto, ON M9P 3T9
416 614.6769

FOR MEDICAL EMERGENCIES AND TRANSPORTATION EMERGENCIES SUCH AS SPILLS,
LEAKS OR FIRES, CALL 1-800-424-9300 (24 HOURS).

NET CONTENTS: 5 X 16 g

DO NOT SELL THE SOLUBLE PACKS SEPARATELY.

DO NOT USE A KNIFE OR OTHER SHARP OBJECT TO OPEN THIS OVERPACK

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN. Harmful if inhaled or absorbed through the skin. Avoid breathing vapors or spray mist. Avoid contact with eyes, skin or clothing. Keep out of reach of domestic animals. During all mixing/loading and clean-up and repair activities, and when applying to outdoor ornamentals, wear long sleeved shirt, long pants and chemical resistant gloves. In greenhouses wear a chemical-resistant rain suit, chemical resistant gloves, rubber boots, goggles or face shield and either a respirator with a NIOSH/MSHA/BHSE-approved organic vapour-removing cartridge with a pre-filter approved for pesticides or a NIOSH/MSHA/BHSE-approved canister approved for pesticides during application. Workers should not enter treated areas for 12 hours following application. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining personal protective equipment. If no such instructions for washables, use detergent and hot water. Keep and wash personal protective equipment separately from other laundry. Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove personal protective equipment immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL PRECAUTIONS:

Do not apply TRISTAR 70 WSP Insecticide directly to water. Do not contaminate water used for irrigation or domestic purposes. Do not contaminate aquatic habitats such as sloughs, coulees, ponds, prairie potholes, lakes, rivers, streams, reservoirs and wetlands or terrestrial habitats such as forested areas, shelter belts, wood lots, hedgerows, pastures, range lands and shrub lands when cleaning and rinsing spray equipment or containers. Do not allow effluent or runoff from greenhouses containing this product to enter lakes, streams, ponds, or other waters.

Overspray or drift to these sensitive habitats should be avoided.

A buffer zone of 20 metres for application by ground boom sprayer and a buffer zone of 30 metres for application by air-blast/vineyard sprayer is required between the downwind point of direct application and the closest edge of sensitive aquatic habitats including sloughs, coulees, ponds, prairie potholes, lakes, rivers, streams, reservoirs and wetlands.

A buffer zone of 2 metres for application by ground boom sprayer and a buffer zone of 10 metres for application by air-blast/vineyard sprayer is required between the downwind point of direct application and the closest edge of sensitive terrestrial habitats such as forested areas, shelter belts, wood lots, hedgerows, pastures, range lands and shrub lands.

For ground application, do not apply during periods of dead calm or when winds are gusty.

For Air-blast/vineyard sprayer, do not direct spray above trees/vines and turn off outward pointing nozzles at row ends and outer rows. Do not apply during periods of dead calm or when winds are gusty or when wind speed is greater than 16 km/hr at the application site as measured outside of the orchard/vineyard on the upwind side.

Acetamiprid is toxic to honey bees exposed to direct treatment. Do not apply when bees are present in the area to be treated.

HANDLING PRECAUTIONS:

Do not handle the inner bag with wet gloves as the bag will begin to dissolve on contact with even a small amount of water. Do not open the inner bags or attempt to remove the product from them. The water soluble packet is not designed to be re-measured or subdivided. The inner bag should not be opened and partial bag not used. The entire contents of the inner (water soluble) bag must be used.

FIRST AID:

Contact a physician IMMEDIATELY in all cases of suspected poisoning.

IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take the container label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION:

There is no specific antidote. All treatment should be based on observed signs and symptoms of distress in the patient. Overexposure to materials other than this product may have occurred.

STORAGE CONDITIONS:

Do not store in or around the home. Store unused product in a cool, ventilated, dry, locked area. Do not allow prolonged storage in areas where temperatures frequently exceed 46°C (115°F). Never transfer this product to another container for storage.

DISPOSAL:

1. Make the empty container unsuitable for further use.
2. Dispose of the container in accordance with provincial requirements.
3. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill and for clean-up of spills.

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

TRISTAR 70 WSP Insecticide

Contains Acetamiprid

AGRICULTURAL

GUARANTEE: Acetamiprid.....70.35% by weight



READ THE LABEL AND BOOKLET BEFORE USING

KEEP OUT OF REACH OF CHILDREN

REGISTRATION NO. 27127 PEST CONTROL PRODUCTS ACT

Registered by:

Nippon Soda Co., Ltd.
2-1, 2-Chome Ohtemachi
Chiyoda-ku, Tokyo 100-8165,
Japan

Agent: David Latter
D. W. Latter and Associates
44 Tamarack Circle
Toronto, ON M9P 3T9
416 614.6769

FOR MEDICAL EMERGENCIES AND TRANSPORTATION EMERGENCIES SUCH AS
SPILLS, LEAKS OR FIRES, CALL 1-800-424-9300 (24 HOURS).

NET CONTENTS: 16 g

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN. Harmful if inhaled or absorbed through the skin. Avoid breathing vapors or spray mist. Avoid contact with eyes, skin or clothing. Keep out of reach of domestic animals. During all mixing/loading and clean-up and repair activities, and when applying to outdoor ornamentals, wear long sleeved shirt, long pants and chemical resistant gloves. In greenhouses wear a chemical-resistant rain suit, chemical resistant gloves, rubber boots, goggles or face shield and either a respirator with a NIOSH/MSHA/BHSE-approved organic vapour-removing cartridge with a pre-filter approved for pesticides or a NIOSH/MSHA/BHSE-approved canister approved for pesticides during application. Workers should not enter treated areas for 12 hours following application. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining personal protective equipment. If no such instructions for washables, use detergent and hot water. Keep and wash personal protective equipment separately from other laundry. Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove personal protective equipment immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL PRECAUTIONS:

Do not apply TRISTAR 70 WP Insecticide directly to water. Do not contaminate water used for irrigation or domestic purposes. Do not contaminate aquatic habitats such as sloughs, coulees, ponds, prairie potholes, lakes, rivers, streams, reservoirs and wetlands or terrestrial habitats such as forested areas, shelter belts, wood lots, hedgerows, pastures, range lands and shrub lands when cleaning and rinsing spray equipment or containers. Do not allow effluent or runoff from greenhouses containing this product to enter lakes, streams, ponds, or other waters.

Overspray or drift to these sensitive habitats should be avoided.

A buffer zone of 20 metres for application by ground boom sprayer and a buffer zone of 30 metres for application by air-blast/vineyard sprayer is required between the downwind point of direct application and the closest edge of sensitive aquatic habitats including sloughs, coulees, ponds, prairie potholes, lakes, rivers, streams, reservoirs and wetlands.

A buffer zone of 2 metres for application by ground boom sprayer and a buffer zone of 10 metres for application by air-blast/vineyard sprayer is required between the downwind point of direct application and the closest edge of sensitive terrestrial habitats such as forested areas, shelter belts, wood lots, hedgerows, pastures, range lands and shrub lands. For ground application, do not apply during periods of dead calm or when winds are gusty.

For Air-blast/vineyard sprayer, do not direct spray above trees/vines and turn off outward pointing nozzles at row ends and outer rows. Do not apply during periods of dead calm or

when winds are gusty or when wind speed is greater than 16 km/hr at the application site as measured outside of the orchard/vineyard on the upwind side.

Acetamiprid is toxic to honey bees exposed to direct treatment. Do not apply when bees are present in the area to be treated.

HANDLING PRECAUTIONS:

Do not handle the inner bag with wet gloves as the bag will begin to dissolve on contact with even a small amount of water. Do not open the inner bags or attempt to remove the product from them. The water soluble packet is not designed to be re-measured or subdivided. The inner bag should not be opened and partial bag not used. The entire contents of the inner (water soluble) bag must be used.

FIRST AID:

Contact a physician IMMEDIATELY in all cases of suspected poisoning.

IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take the container label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION:

There is no specific antidote. All treatment should be based on observed signs and symptoms of distress in the patient. Overexposure to materials other than this product may have occurred.

STORAGE CONDITIONS:

Do not store in or around the home. Store unused product in a cool, ventilated, dry, locked area. Do not allow prolonged storage in areas where temperatures frequently exceed 46°C (115°F). Never transfer this product to another container for storage.

DISPOSAL:

1. Make the empty container unsuitable for further use.
2. Dispose of the container in accordance with provincial requirements.
3. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill and for clean-up of spills.

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

DIRECTIONS FOR USE:

FOR ORNAMENTAL USE ONLY. NOT FOR WOODLANDS OR FOREST MANAGEMENT.

FOR USE ON ORNAMENTAL AND FLOWERING PLANTS GROWN OUTDOORS AND IN GREENHOUSES, SHADEHOUSES AND LATHHOUSES: Bedding plants, flowers grown for cuttings, foliage plants, potted flowering plants, ornamental trees, and non-bearing fruit and nut trees. (Non-bearing crops are perennial crops that will not produce a harvestable raw agricultural commodity during the season of application). Do not apply to bearing fruit trees.-

For control of ornamental pests, apply broadcast spray thoroughly and uniformly to all exposed tissues. To obtain thorough and uniform spray coverage, choose a finished spray volume appropriate for the size of the plants and the amount of foliage which will provide thorough coverage throughout the canopy. For optimum control, allow at least 6 hours before overhead irrigation of foliage. Do not allow public use of treated area during application.

Mix TriStar 70 WSP Insecticide with sufficient water to obtain thorough and uniform foliar spray coverage. Apply as soon as insects reach treatment thresholds. See resistance management section if multiple sprays are needed.

Do not reapply more than once every 7 days. To determine whether application or re-application is necessary, monitor pest densities and consult provincial guidelines and/or local extension experts.

Note: Since ornamental varieties are numerous and constantly changing and may react differently at various sites, test the product on a small scale before making large-scale applications if there is not local experience.

PESTS	RATE WATER SOLUBLE PACKS/1000 L	COMMENTS
Aphids, European pine sawfly	3	Apply as a full coverage foliar spray. One water soluble pack per 380 L is adequate for aphid control in most conditions.
Tentiform leaf miner, Leafhoppers	5	
Whiteflies	5-10	

Product can be applied with the water volume needed to provide thorough coverage.
If applying less than 1000L/ha, consider recommended rates as rate per hectare.
Use the higher rate when insect pressure is high.

USE RESTRICTIONS AND PRECAUTIONS:

- For ornamental use only. Not for woodlands or forest management use.
- Do not make more than 2 applications per year per greenhouse, shadehouse or lathhouse.
- For outdoor ornamentals, do not make more than 5 applications per year.
- Do not reapply more than once every 7 days.
- Do not apply more than 55 water soluble packs/ha (616 g ai/ha) per year.
- Do not apply to bearing fruit trees or vegetables.

Greenhouse Peppers: Control of aphids

NOTE TO BUYER/USER: READ THE FOLLOWING BEFORE USING THIS PRODUCT FOR SPECIAL USE APPLICATIONS:

The **DIRECTIONS FOR USE** for this product for the uses described on the label were developed by persons other than Nippon Soda Co., Ltd. and accepted for registration by Health Canada under the User Requested Minor Use Registration Label Expansion program. Nippon Soda Co., Ltd. itself makes no representation or warranty with respect to performance (efficacy) and/or crop tolerance (phytotoxicity) claims for this product when used in accordance with this label.

Accordingly, the Buyer and User assumes risk of damage or loss resulting from such uses, and agrees to hold Nippon Soda Co., Ltd. harmless from any claims based on efficacy and/or phytotoxicity in connection with the uses described on this label.

To determine whether application or re-application is necessary, monitor pest densities or, where available, consult provincial guidelines and/or local extension experts. To manage insect resistance, alternate applications with an insecticide with a different mode of action.

Greenhouse Peppers: Control of aphids

Note: Since pepper varieties are numerous and may react differently at various sites, test the product on a small scale before making large-scale applications if there is not local experience.

Mix TRISTAR 70 WSP Insecticide with sufficient water to obtain thorough and uniform coverage of the foliage. Use a spray volume appropriate for the size of the plants being treated. Apply when aphid numbers have exceeded the action threshold as indicated by crop monitoring. For optimum control, allow at least 6 hours before overhead irrigation.

PESTS	RATE and COMMENTS
Aphids	Apply in a spray concentration of 3 water soluble packs/1000 L of water. Do not exceed 5 water soluble packs (1667 L of spray solution) per hectare.

USE RESTRICTIONS AND PRECAUTIONS

- For use on peppers grown in greenhouses only.
- Do not make more than 2 applications per year per greenhouse.
- Do not reapply more than once every 7 days.
- Pre-harvest Interval: Do not apply within 3 days of harvest.

FOR SALE FOR USE FOR THE CONTROL OF WHITEFLIES ON

GREENHOUSE TOMATOES:			
SITE	PEST	RATE WATER SOLUBLE PACKS	SPECIFIC DIRECTIONS
Greenhouse tomatoes	Whiteflies	15 packs per 2 ha (1 pack /1333 m ²)	Apply in sufficient water to provide adequate distribution to the treated area.
<p>Do not make more than 2 applications per year per greenhouse. Apply when whitefly numbers have exceeded the action threshold as indicated by crop monitoring. Apply through drip irrigation to the growing media. Drip irrigation is the only application method approved. Pre-harvest interval is 1 day. To determine whether application or re-application is necessary, monitor past densities and consult provincial guidelines and/or local extension experts. Alternate applications with an insecticide with a different mode of action to manage insect resistance. Repeat if necessary at a 21-day treatment interval.</p>			

RESISTANCE MANAGEMENT

For resistance management, this is a Group 4 insecticide. Any insect population may contain individuals naturally resistant to this product and other Group 4 insecticides. The resistant biotypes may dominate the insect population if these insecticides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism,, may also exist. Appropriate resistance management strategies should be followed.

To delay insecticide resistance:

- Where possible, rotate the use of this product with different groups that control the same pests.
- Use tank mixtures with insecticides from a different group when such use is permitted.
- Insecticide use should be based on an integrated pest management (IPM) program that includes scouting and record keeping and considers cultural, biological, and other chemical control practices;
- Monitor treated pest populations for resistance development.
- Contact the local extension specialist or certified crop advisors for any additional pesticide resistance-management and IPM recommendations for the specific site and pest problems in the area.
- For further information or to report suspected resistance, contact Nippon Soda Co., Inc. at 1-212-490-0350

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.