

19-JAN-2007

SePRO

**A-Rest
Solution**

- **For Professional Use**
- **Horticultural**

GUARANTEE: Ancyimidol..... 264 mg/L

CONTAINS BENZOIC ACID AT 0.0002% AND
1-(3-CHLOROALLYL)-3,5,7-TRIAZA-1-AZONIAADAMANTANE
CHLORIDE AT 0.13% AS PRESERVATIVES.

REGISTRATION NO. 16393 PEST CONTROL PRODUCTS ACT

COMMERCIAL

NET CONTENTS: 113.6 L

READ THE LABEL BEFORE USING

CAUTION:

KEEP OUT OF REACH OF CHILDREN. MAY BE HARMFUL IF
SWALLOWED. AVOID CONTACT WITH SKIN, EYES, OR
CLOTHING. AVOID CONTAMINATION OF FOODSTUFFS, FEEDS
AND FISH PONDS.

Distributed by:

Plant Products Co. Ltd.
314 Orenda Road
Brampton, Ontario L6T 1G1

Manufactured by:

SePRO Corporation
11550 North Meridian Street, Suite 600
Carmel, Indiana, U.S.A. 46032

Telephone Number :
(317) 580-8282

Notification Change

FIRST AID

Take container, label or product name and Pest Control Product Registration Number with you, when seeking medical attention.

If swallowed: Do not induce vomiting. Call a physician or Poison Control Centre.

If in eyes: Flush eyes with plenty of water. Get medical attention or contact a poison control centre if irritation persists.

If on skin: Wash with plenty of soap and water. Get medical attention or contact a poison control centre if irritation develops.

If inhaled: Remove individual to fresh air. If breathing difficulty occurs, get medical attention or contact a poison control centre.

TOXICOLOGICAL INFORMATION: Treat symptomatically.

PRECAUTIONS:

Wear long pants, a long-sleeved shirt, chemical-resistant gloves and boots during mixing/loading, application, clean-up and repair activities.

Do not re-enter or allow worker entry into treated areas until 12 hours after application.

ENVIRONMENTAL PRECAUTIONS:

Do not allow waste water, effluent or runoff containing this product to enter lakes, streams, ponds or other waters.

GENERAL INFORMATION

Read all directions carefully before applying.

A-REST for Height Control of Container-Grown Lilies, Poinsettias, and Chrysanthemum A-REST is a plant growth regulator, which effectively reduces internode elongation resulting in more desirable compact plants. When used as directed, A-REST produces no phytotoxic effects.

A-REST is not a substitute for any cultural practices (lighting, watering and temperature control). Use optimum cultural practices for growing plants listed on this label.

APPLICATION DIRECTIONS

Mix A-REST with water during spray tank filling operation. Specific rates will vary with plant species, variety, timing and type of treatment, and growing conditions. The degree of response desired often varies with location and individual cultural practices. The rate of application within the recommended range depends upon the degree of growth inhibition desired by the grower and should be established by using the product on a limited basis at varying rates.

The amount of growth inhibition from A-REST treatment depends upon the total quantity of active ingredients applied per plant. Treatment techniques and equipment should ensure uniform distribution of A-REST within the treated area.

SOIL DRENCH APPLICATIONS

Chrysanthemums

Dilute the A-REST solution as indicated in Table I, and apply as a drench treatment to moisten soil. Treat plants 5 - 15 cm in height (approximately two weeks following pinch).

Lilies

Dilute the A-REST solution as indicated in Table I, and apply as a drench treatment to moisten soil. Treat plants from emergence to 30 cm in height. For optimum results, treatments should be made to plants 5 to 15 cm tall.

Poinsettias

Dilute the A-REST solution as indicated in Table I, and apply as a drench treatment to moisten soil. Treat plants at pinch to 4 weeks after pinch, or 8 to 12 weeks before finishing.

TABLE I

No. of 15-cm Pots to be Treated	Drench Volume (mL)	Dilution Ratio A-REST to Water	Rate (mg/15-cm Pot)	Final Drench Concentration (mg/L)
250	120	250 mL 29.75 L	0.25	2
250	60	250 mL 14.75 L	0.25	4
250	60	500 mL 14.5 L	0.50	8
250	120	500 mL 29.5 L	0.50	4

WARNING

Do not add any additional wetting agents to A-REST solutions. Plants treated with A-REST may require less water than untreated plants. Over-application or uneven application of A-REST will result in excessive or irregular growth control. A slight delay (two to five days) in flower development may occur at the higher recommended treatment levels.

DO NOT FREEZE

DISPOSAL

Returnable Containers

Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the Pest Control Products Act to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.